



# Attachment style and family presence preference during invasive nursing procedures

Nurten Kaya<sup>1</sup>, Nuray Turan<sup>2\*</sup>

<sup>1</sup>Health Sciences Faculty, Istanbul University Cerrahpasa, Istanbul, Turkey, <sup>2</sup>Florence Nightingale Nursing Faculty, Istanbul University Cerrahpasa, Istanbul, Turkey

## ABSTRACT

**Introduction:** The attachment style and family presence preference are important during the invasive medical procedures. Our aim was to analyze the effects of adult attachment styles of the patients which prefer presence of their family members during the invasive medical procedures in emergency departments.

**Methods:** We included 76 randomly selected patients who received the invasive medical procedures in the emergency department of the University hospital. The Patient Information Form and Relationship Scales Questionnaire were used to collect data.

**Results:** About 57.9% of the patients said that they preferred their relatives to stand by them during invasive nursing procedures. 56.6% of participants stated that they favor their relatives to support them at the time of such interventions. Average scores of adult attachment styles were  $3.02 \pm 0.63$  for fearful,  $3.57 \pm 0.57$  for dismissing,  $2.87 \pm 0.50$  for preoccupied, and  $2.79 \pm 0.66$  for secure attachment style. Adult attachment styles of participants were found to have no impact on preferring someone standing by them at the time of invasive nursing interventions ( $p > 0.05$ ).

**Conclusion:** Adult attachment styles do not affect the patients' need to have a family member stand beside them during an invasive medical procedure.

**Key words:** Attachment style; emergency department; family presence; invasive nursing procedures

## INTRODUCTION

Emergency departments offer life-saving and therapeutic treatment responses. Thus, it is vital to have cooperative synergy and awareness within the team

and it is required that the person as well as his family is assessed in an integrated manner. During the emergency hospitalization of patients, some unfavorable consequences can be sometimes seen as a result of medical care operations and interventions exercised. The ethical approach of making no harm is taken into account in due form in the exercise of such interventions, but it is needed that nurses endeavor more to decrease the negative impact of applied intervention. However, there have been few studies conducted on the presence of families at the time of invasive nursing interventions (1-3).

\*Corresponding author: Nuray Turan, Department of Fundamentals of Nursing, Florence Nightingale Nursing Faculty, Istanbul University, Cerrahpasa, bide-i Hurriyet Street, 34381 Sisli/Istanbul, Turkey.  
E-mail: [nkaraman@istanbul.edu.tr](mailto:nkaraman@istanbul.edu.tr)

Submitted: 01 September 2018/Accepted: 09 December 2018

DOI: <https://doi.org/10.17532/jhsci.2018.188>



Family presence refers to the availability of the family in the spot so that they are able to contact the patient visually or physically at the time of invasive interventions and resuscitation in medical centers (4,5). A good deal of qualified medical institutions encourages the presence of family at the time of diagnosis, care, and treatment processes. They have issued directives related to the presence of family at the time of such procedures. The Emergency Nurses Association was the first to introduce such directive in 1994 and revise it in October 2005 with a view to approve family presence (6).

An integrative medical care incorporates patient as well as family (7,8). In pediatric literature, the presence of family during invasive interventions is usually underlined, but various studies conducted on other units that offer medical care to people from various ages have dealt with this issue as well (9).

Nevertheless, attachment is a sentimental connection between two people, and as part of this connection, it is expected that one or both of these people will offer care and shelter when necessary. Behaviors of people who are related to each other are influenced by the type of attachment. Attachment styles are influential on the behaviors of the individuals who are in relationships with one another; and also influential in obtaining satisfaction from their relationships, in the level with which the individuals are affected by the problems experienced in relationships, and in coping with these problems (10-12).

Still, the theory of attachment gives pedagogues, clinicians, and academicians a means to investigate the effect of past livings on later adaptation as it is useful in explaining personal differences in terms of emotional regulation, stress reactions, and interpersonal behavior. The attachment theory states that experiences that people have early with caregivers are turned into internal intellectual manifestations of attachment throughout the adulthood (13-16).

Bartholomew and Horowitz (10) have developed an adult attachment model consisting of four categories and it is partake of Bowlby's original models on the self and others. Bartholomew and Horowitz (10) suggest that adult people with a positive model of other individuals as present and caring and themselves

as deserving approval and care may be identified as securely attached. It is considered that people who are securely attached had a stable, caring, and conscious caregiving history and tend to establish aiding relationships successfully. Hence, these people conveniently use other individuals as a means of aid when necessary (12).

The other three types of attachment include a negative working model regarding the self or others, and therefore, they are all called "insecure" types of attachment. People who have insecure attachment style are inclined to be more concerned and show higher level of self-protective approach to avoid possibly supportive affairs. People who have "fearful" insecure attachment are tend to not trust others, avoid any affairs, and consider themselves as unpleasant and not worthy of aid. These "fearful" people may seem like to be driven by closeness with other people since it can make up for their perceived unfavorable image, and they escape from close affairs due to fear of rejection (10).

People who have a positive model toward others and a negative one toward themselves are considered as abstracted in insecure types of attachment. They are concerned with the requirements for attachment, and thus, they determinedly try to be accepted and approved by means of attached relationships. These preoccupied attachments are inclined to sentimentally unbalanced, with little self-esteem and insistent will to get approval (11).

On the other hand, dismissing style of attachment represents a positive model of the self but negative model of others. Individuals with such style are inclined to be very doubtful close affairs and have fear of letting themselves to trust others (12,17).

Our aim was to examine the effect of adult attachment styles regarding the fact that the patients prefer their family members or relatives or the people they care to stand by them during invasive nursing procedures.

## METHODS

### Research questions

- What are the demographic and disease variables of patients?

- What does the patient think about family presence preference at the time of invasive nursing procedures?
- What is the attachment style of patients?
- Do the attachment style of patients have an impact on their family presence preferences?

### Study population and sample

This study was conducted in the observation unit of the internal medicine section located in the emergency department of a university hospital in Istanbul. The sample consisted of 76 patients who were selected by random sampling method. The following inclusion criteria were used:

- Being older than 18,
- Being open to communication and cooperation,
- Undergoing an invasive nursing procedure during hospitalization in the emergency unit, and
- Having given informed consent for participation in the study.

Those who developed complications during the intervention or were unaccompanied were excluded from the study.

### Instruments

The Patient Information Form and the Relationship Scales Questionnaire (RSQ) were used to collect data.

#### *The patient information form*

It was created based on the existing literature (18,19) and information gathered on sociodemographic variables such as gender, age, marital status, living arrangements, educational status, occupation, and medical diagnosis. Questions were found which are given below:

- Would you like your family with you during nursing interventions?
- If your answer is yes, why you would?
- If your answer is no, why you wouldn't?
- Is there anyone, especially you want during the process?
- If your answer is yes, who is this person?
- Who should decide that someone have to stay with you or not, during the interventional nursing practices? (Options: Me, my familiar, the nurse, doctor, etc., more than one response can be given.)

#### *The RSQ*

The questionnaire was utilized to find out attachment styles of patients and created by Griffin and Bartholomew (16). In 1999, Sumer and Gungor determined the validity and reliability of its Turkish version (20). The RSQ consists of 17 items, which are used to form continuous subscales of attachment style categories. These items include five-point Likert scales. These scales are used to measure the level of statements in each item to constitute respondent's opinions about close interpersonal relationships. Regarding the relationship style domains, there are four attachment style domains, which are secure, fearful, preoccupied, and dismissing. The continuous scores represent such attachment styles and are gathered with questions intended to measure such styles and by means of division of this total score by the number of questions in each subscale. Therefore, the scores obtained from these subscales may differ from 1 to 5. To categorize the participants based on their attachment styles, continuous scores obtained through this method are utilized. In this grouping process, participants are allocated to the attachment group where they get the highest score (20).

### Ethical consideration

A written consent was obtained after submitting an information form including the aim and scope of the study. The participants making up the sample were informed about its aim and benefits apart from their roles in the study under the principles of willingness and volunteerism, and their consent was gained.

### Statistical analysis

To statistically analyze 76 participants, Statistical Package for the Social Sciences 22.0 for Windows program was used. Moreover, descriptive statistics (i.e., frequency, percentages, means, and standard deviations) were also determined. For the analysis of categorical data, Pearson's Chi-square and Fisher's exact tests were used. The independent samples t-test was used to compare the parametric data. The results were assessed at a confidence interval of 95%, and significance threshold for primary analyses was set at 0.05.

## RESULTS

Table 1 shows the demographic and disease features of the participants. Accordingly, 56.6% of the participants were female, their average age was  $47.83 \pm 18.88$ , 65.8% of them were married, 65.8% of them had children, 56.6% of them were living with their families, 28.9% of them had university degree or higher education degree, 36.8% of them were state officials, 93.4% of them had general health insurance, and the other 6.6% of them had green cards.

Table 2 shows the opinions of participants regarding their family presence preference at the time invasive nursing interventions. Accordingly, 56.6% of participants stated that they favor their relatives to support them at the time of such interventions. 51.3% of patients said yes in response to the question of “Is there someone you would like him/her to stand by your especially during the intervention?” Regarding the persons they prefer to be with them during invasive nursing procedures, the participants said that they prefer their mother/father, child, spouse, sibling, relatives, and friends. 76.3% of participants answered the question of “who should decide that someone have to stay with you or not, during the interventional nursing practices?” as “me.”

Regarding the mean point of adult attachment styles of participants of the study, it was found that potential point distribution is between 1 and 5 in all styles, 3.57 in dismissing,  $3.02 \pm 0.63$  in fearful,  $2.87 \pm 0.50$  in preoccupied, and  $2.79 \pm 0.66$  in secure (Table 3).

Adult attachment styles of participants were found to have no impact on the condition of preferring someone standing by them at the time of invasive nursing interventions ( $p > 0.05$ ; Table 4).

## DISCUSSION

According to a study, anxious as well as avoidant aspects of attachment are related to health risk behaviors (21).

Any association of curative connections with the patient's agreeability or choice to establish relationships has not been proven, which indicates that therapeutic relationship and service attachment measurements assess a dimension that is noticeable regarding the experience of patients.

**TABLE 1.** Distribution of demographic and disease variables of the patients ( $n=76$ )

Demographic and disease variables	<i>n</i> (%)
Gender	
Female	43 (56.6)
Male	33 (43.4)
Age categories	
18–31	26 (34.2)
32–45	8 (10.5)
46–59	18 (23.7)
60–↑	24 (31.6)
Age (Minimum–Maximum)	(19–89)
Mean±SD	$47.83 \pm 18.88$
Marital status	
Married	50 (65.8)
Single/Widowed/Divorced	26 (34.2)
Having children	
Yes	50 (65.8)
No	26 (34.2)
Lived with	
Alone	9 (11.8)
Family	43 (56.6)
Only spouse	15 (19.7)
Relative	4 (5.3)
Friend	5 (6.6)
Educational status	
Illiterate	9 (11.8)
Literate	10 (13.2)
Primary school	14 (18.4)
Secondary school	21 (27.6)
Higher education and above	22 (28.9)
Occupation	
Not working	23 (30.3)
State official	28 (36.8)
Worker	6 (7.9)
Freelancer	5 (6.6)
Retired	14 (18.4)
Social security	
General health insurance	71 (93.4)
Green cards	5 (6.6)
Medical diagnosis	
Rheumatic diseases	7 (9.2)
Endocrinologic diseases	9 (11.8)
Hematologic diseases	21 (27.6)
Cardiological diseases	7 (9.2)
Infectious diseases	9 (11.8)
Nephrologic diseases	16 (21.1)
Lung diseases	5 (6.6)
Neurologic diseases	2 (2.6)

SD: Standard deviation

**TABLE 2.** Distribution of thought about the patients' family presence preference during invasive nursing procedures

Thoughts about patients' family presence preference during invasive nursing procedures	n (%)
Requesting family during the invasive nursing procedures	
Yes	43 (56.6)
No	33 (43.4)
The reason why family is requested	
The patients who do not want their family (preference is no)	33 (43.4)
They become a support and help to me	20 (26.3)
I feel comfortable, and safe and makes me	10 (13.3)
My pain decreases	1 (1.3)
My fear/anxiety decreases	8 (10.5)
They witness to procedures	2 (2.6)
No reason	2 (2.6)
The reason why family is not requested	
The patients who want their family (preference is yes)	43 (56.6)
It does not matter/not necessary	19 (25.0)
My family worries/gets upset	5 (6.6)
I do not want make them see me during procedure/I get angry	8 (10.5)
I trust nurses	1 (1.3)
The presence of the person who is significantly requested by patient during procedure	
Yes	39 (51.3)
No	37 (48.7)
The person who is significantly requested by the patient during procedure	
Who responses no	37 (48.7)
The parents	5 (6.6)
His/her children/child	9 (11.9)
Spouse	20 (26.3)
Brother/sister	3 (3.9)
Relative/friend	2 (2.6)
The person who prefers who should stay with the patient during procedure*	
Me (the patient)	58 (76.3)
The family	26 (34.2)
The nurse	15 (19.7)
The doctor	8 (10.5)

\*Multiple responses are given

Accordingly, measurement of team attachment and therapeutic relationship is corresponding to measurement of almost identical structures. The service users who have higher level of preoccupied attachment styles are likely to experience more frustration in establishing a positive attachment to departments that are modified again and again (22).

The rates of prevalence for secure, dismissing, preoccupied, and fearful attachment styles in 4095 primary care diabetes patients were found to be 44.2%, 35.8%, 7.9%, and 12.1%, respectively. In

a comparison of secure attachment style, dismissing attachment style was found to be related with very low level of physical activity, foot care, diet, and conforming to oral hypoglycemic drugs and high level of smoking, and patient-provider relationship was used to intervene in such associations. It was also found that preoccupied attachment style which features overreliance on other people is correlated with very low risk of glycosylated hemoglobin rates >8%, by comparison with secure attachment. Therefore, self-management and outcomes of diabetes are

related with the attachment style to a significant extent (23).

Although medical care is used in interpersonal settings, the knowledge about the relevance of interpersonal choices or patterns of service users is limited. A few studies have been conducted to discover the connections between attachments, which is pattern of reliance on others and the style of benefiting from medical care.

In this report, examined how attachment characteristics predicted the frequency of digital rectal exam and prostate-specific antigen testing in a sample of African-descent men. 414 African-descent men who are aged between 45 and 70 performed measures of prostate screening and attachment as well as predictors of screening (demographics, insurance, family history, physician variables, knowledge, perceived risk, and accessibility). In line with the estimations, dismissiveness, which is seen most frequently as a relational pattern in the elderly, anticipated frequent prostate-specific antigen testing and digital rectal examination, while attachment security, which refers to contentment about close affairs, anticipated

lower screening frequency. Identifying the interpersonal characteristics predicting screening may help identify men at risk of suboptimal health care use and guide the development of interventions suited to the normative relational preferences of current cohorts of older, African-descent men (24).

We found associations between psychological attachment anxiety on smoking and higher number of session use, independent of disease severity, which was more pronounced for women. After diagnosis of CP and AgP, individuals who have more attachment avoidance opted for periodontal treatment afterward. In terms of attachment avoidance and anxiety and number of teeth at the initial stage of treatment, differential connections were identified in men. Besides recognized psychosocial risk factors, psychological attachment styles can be used to explain periodontal disease (25).

In this study, the function that attachment insecurity has in cervical screening behaviors and obstacles in a group of 257 female undergraduates was examined. Information related to aspects of attachment and attachment patterns was gathered. Attachment anxiety and attachment avoidance were associated with decreased likelihood of having participated in cervical screening and positively associated with screening barriers. Furthermore, it was found that people who have attachment insecurity (preoccupied, fearful, and dismissing) tend to have higher level of screening obstacles and dismissing individuals avoided being involved in screening when compared with secure individuals. These findings prove that insecure attachment could pose a risk for

**TABLE 3.** The patients' attachment style scores according to RSQ\*

Attachment styles	Minimum–Maximum	Means±SD
Fearful	1.50–4.25	3.02±0.63
Dismissing	2.40–4.80	3.57±0.57
Preoccupied	2.00–4.50	2.87±0.50
Secure	1.40–4.60	2.79±0.66

\*RSQ: The Relationship Scales Questionnaire: Range 1–5. SD: Standard deviation

**TABLE 4.** The patients' family presence preference according to attachment styles scores

Attachment styles	Family presence preference	Means±SD	t	p
Fearful	Yes	3.11±0.65	1.532	0.130
	No	2.89±0.60		
Dismissing	Yes	3.52±0.58	-0.816	0.417
	No	3.63±0.57		
Preoccupied	Yes	2.93±0.50	1.244	0.217
	No	2.79±0.48		
Secure	Yes	2.71±0.75	-1.260	0.212
	No	2.91±0.51		

\* Range 1–5. SD: Standard deviation



insufficient cervical screening and screening obstacles (26).

It was found that there is an important difference when it comes to being informed of surroundings for attachment patterns. Furthermore, a significant association was determined between attachment styles of patients (fearful, preoccupied, and dismissing) and intensive care experience. At the time of discharge from intensive care unit, secure attachment style points and Acute Physiology and Chronic Health Evaluation II score of the patients were found to be significantly associated. Therefore, in the course of planning and exercising nursing care and procedures for people in an intensive care unit, it is important to take note of attachment patterns (3).

According to the study, attachment to the caregiver could be a significant predictor of the therapeutic alliance. Based on the findings, attachment toward the caregiver could be of more significance compared to prior manifestations of attachment in the context of therapeutic alliance (27).

Participants who have medically unidentified symptoms generally exhibit insecure adult attachment styles, and for this reason, more attendance records could be classified as pathological behavior looking for care, which is associated with insecure attachment. Caregivers could handle patients in a different manner by considering constant visit as a pathological act shown with a view to get help (28).

Typical styles in terms of close affairs, sentimental regulation, having social support, behavioral descriptors, metallization, and narrative coherence characterize the four adult attachment styles (secure, preoccupied, dismissing, and fearful). They are associated with medical care affiliations and their results. Typical narrations of adult attachment styles could help doctors understand personal differences in interpersonal structure that affect the health of participants (29).

Attachment styles of patients could be a significant determinant in their demands for physician-assisted dying. Furthermore, a physician's capability to have a productive relationship with his patient during the course of dying can be improved by understanding his attachment style (30).

According to the findings, there is a link between patients who have a hair loss that is not visible and the contradictory attachment style. Overall clinical

effects and attachment indicators, including attachment security and the methods to deal with, have been found to be an important input to anticipate changes in the quality of life scales, "self-esteem" and "emotions." Thus, attachment security is likely to be a basic instrument that reconciles with subjective health and the patients who have female alopecia are likely to have special attachment vulnerability. Future studies are needed to elaborate on the pertinence of attachment styles in doctor and patient relationship and on psychotherapeutic procedures (31).

### Limitations

The current study was conducted on emergency department of a university hospital in Istanbul. In the context of our research, there had been a limited number of samples. Therefore, the results cannot be generalized. Further longitudinal research is, therefore, recommended to gather data from health-care members in the emergency department that can be compared and contrasted to explore changes in attachment and family presence preference.

### CONCLUSION

In empirical terms, this study suggests that attachment pattern of patients had no impact on their family presence preference at the time of invasive nursing interventions. Based on these findings, it is possible to recommend the following:

- The patients themselves should choose family support at the time of invasive nursing interventions. It is better to have the family stand by the patient in case no obstacle exists at the time of such procedures. However, it is better to have the family outside depending on the preference of patient if he is not willing to have the family around him in the course of these procedures.
- Different departments should be used in future studies conducted on this topic.
- It is needed to conduct qualitative studies regarding the family presence at the time of interventional nursing procedures.

### FUNDING

We present our thanks to Istanbul University Scientific Research Projects Unit for their contributions to our study (BEK-2017-25520).

## REFERENCES

- İşlekdemir B, Kaya N. Effect of family presence on pain and anxiety during invasive nursing procedures in an emergency department: A randomized controlled experimental study. *Int Emerg Nurs* 2016;24:39-45. <https://doi.org/10.1016/j.ienj.2015.05.008>.
- Kaya N, Kaya H. Effects of attachment styles of nurses on coping strategies. *J Med Sci* 2009;29(6):1563-72.
- Kaya N. Effect of attachment styles of individuals discharged from an intensive care unit on intensive care experience. *J Crit Care* 2012;27(1):103.e7-14. <https://doi.org/10.1016/j.jcrrc.2011.05.020>.
- Emergency Nurses Association. Family Presence at the Bedside During Invasive Procedures and Cardiopulmonary Resuscitation 2005; 2012. Available from: <http://www.ena.org/about/position/pdfs/4e6c256b26994e319f66c65748bfbdbf.pdf>. [Last accessed on 2012 Jan 01].
- McGahey-Oakland PR, Lieder HS, Young A, Jefferson LS. Family experiences during resuscitation at a children's hospital emergency department. *J Pediatr Health Care* 2007;21(4):217-25. <https://doi.org/10.1016/j.pedhc.2006.12.001>.
- Baumhover N, Hughes L. Spirituality and support for family presence during invasive procedures and resuscitations in adults. *Am J Crit Care* 2009;18(4):357-66. <https://doi.org/10.4037/ajcc2009759>.
- Kaya N, Bolol N, Ülgen S, İşiçi Ç, Özen S, Akgün G. Quality of life and influencing factors in caregivers of patients with cancer. *Clin Exp Health Sci* 2018; 8: 25-35. <https://doi.org/10.5152/clinexphealthsci.2017.422>.
- Kaya N. Attachment styles of nursing students: A cross-sectional and a longitudinal study. *Nurse Educ Today* 2010;30(7):666-73. <https://doi.org/10.1016/j.nedt.2010.01.001>.
- Boudreaux ED, Francis JL, Loyacano T. Family presence during invasive procedures and resuscitations in the emergency department: A critical review and suggestions for future research. *Ann Emerg Med* 2002;40(2):193-205. <https://doi.org/10.1067/mem.2002.124899>.
- Bartholomew K, Horowitz LM. Attachment styles among young adults: A test of a four-category model. *J Pers Soc Psychol* 1991;61(2):226-44. <https://doi.org/10.1037/0022-3514.61.2.226>.
- Hawkins AC, Howard RA, Oyeboode JR. Stress and coping in hospice nursing staff. The impact of attachment styles. *Psychooncology* 2007;16:563-72. <https://doi.org/10.1002/pon.1064>.
- Tan A, Zimmermann C, Rodin G. Interpersonal processes in palliative care: An attachment perspective on the patient-clinician relationship. *Palliat Med* 2005;19:143-50. <https://doi.org/10.1191/0269216305pm994oa>.
- Bowlby J. Attachment and Loss. Attachment. Vol. 1. New York: Basic Books; 1969.
- Bowlby J. Attachment and Loss. Separation, Anxiety and Anger. Vol. 2. New York: Basic Books; 1973.
- Bowlby J. Attachment and loss. In: Loss: Sadness and Depression. Vol. 3. London: Hogarth Press; 1980.
- Griffin D, Bartholomew K. Models of the self and other: Fundamental dimensions underlying measures of adult attachment. *J Pers Soc Psychol* 1994;67(3):430-45. <https://doi.org/10.1037/0022-3514.67.3.430>.
- Holmes J. Social relationships: The nature and function of relational schemas. *Eur J Soc Psychol* 2000;30:447-95. [https://doi.org/10.1002/1099-0992\(200007/08\)30:4<447::AID-EJSP10>3.0.CO;2-Q](https://doi.org/10.1002/1099-0992(200007/08)30:4<447::AID-EJSP10>3.0.CO;2-Q).
- Çelik G. Evaluation of patients' families' attitudes to witnessing invasive procedures in the emergency department. *J Acad Emerg Med* 2013;12:63-4. <https://doi.org/10.5152/jaem.2013.028>.
- Mortelmans LJ, Van Broeckhoven V, Van Boxtael S, De Cauwer HG, Verfaillie L, Van Hellemond PL, et al. Patients' and relatives' view on witnessed resuscitation in the emergency department: A prospective study. *Eur J Emerg Med* 2010;17(4):203-7. <https://doi.org/10.1097/MEJ.0b013e328331477e>.
- Sumer N, Gungor D. Psychometric evaluation of adult attachment measures on Turkish samples and a cross-cultural comparison. *Turk J Psychol* 1999;14(43):71-106.
- Ahrens KR, Ciechanowski P, Katon W. Associations between adult attachment style and health risk behaviors in an adult female primary care population. *J Psychosom Res* 2012;72(5):364-70. <https://doi.org/10.1016/j.jpsychores.2012.02.002>.
- Catty J, Cowan N, Poole Z, Ellis G, Geyer C, Lissouba P, et al. Attachment to the clinical team and its association with therapeutic relationships, social networks, and clinical well-being. *Psychol Psychother* 2012;85(1):17-35. <https://doi.org/10.1111/j.2044-8341.2010.02011.x>.
- Ciechanowski P, Russo J, Katon W, Von Korff M, Ludman E, Lin E, et al. Influence of patient attachment style on self-care and outcomes in diabetes. *Psychosom Med* 2004;66(5):720-8. <https://doi.org/10.1097/01.psy.0000138125.59122.23>.
- Conedine NS, Tuck NL, Fiori KL. Attachment and health care utilization among middle-aged and older African-descent men: Dismissiveness predicts less frequent digital rectal examination and prostate-specific antigen screening. *Am J Mens Health* 2013;7(5):382-93. <https://doi.org/10.1177/1557988312474838>.
- Graetz C, Ehrental JC, Senf D, Semar K, Herzog W, Dörfer CE, et al. Influence of psychological attachment patterns on periodontal disease—a pilot study with 310 compliant patients. *J Clin Periodontol* 2013;40(12):1087-94. <https://doi.org/10.1111/jcpe.12159>.
- Hill EM, Gick ML. Attachment and barriers to cervical screening. *J Health Psychol* 2013;18(5):648-57. <https://doi.org/10.1177/1359105312454910>.
- Taylor PJ, Rietzschel J, Danquah A, Berry K. The role of attachment style, attachment to therapist, and working alliance in response to psychological therapy. *Psychol Psychother* 2015;88(3):240-53. <https://doi.org/10.1111/papt.12045>.
- Taylor RE, Marshall T, Mann A, Goldberg DP. Insecure attachment and frequent attendance in primary care: A longitudinal cohort study of medically unexplained symptom presentations in ten UK general practices. *Psychol Med* 2012;42(4):855-64. <https://doi.org/10.1017/S0033291711001589>.
- Maunder RG, Hunter JJ. Assessing patterns of adult attachment in medical patients. *Gen Hosp Psychiatry* 2009;31(2):123-30. <https://doi.org/10.1016/j.genhosppsych.2008.10.007>.
- Oldham RL, Dobscha SK, Goy ER, Ganzini L. Attachment styles of Oregonians who request physician-assisted death. *Palliat Support Care* 2011;9(2):123-8. <https://doi.org/10.1017/S1478951510000660>.
- Schmidt S. Female alopecia: The mediating effect of attachment patterns on changes in subjective health indicators. *Br J Dermatol* 2003;148(6):1205-11. <https://doi.org/10.1046/j.1365-2133.2003.05327.x>.